

# BC Cancer Protocol Summary for Palliative Combination Chemotherapy for Metastatic Colorectal Cancer using Irinotecan, Bevacizumab and Capecitabine

**Protocol Code:**

*GICIRB*

**Tumour Group:**

*Gastrointestinal*

**Contact Physician:**

*GI Systemic Therapy*

## ELIGIBILITY

Patients must have:

- Locally advanced, locally recurrent or metastatic colorectal adenocarcinoma, not curable with surgery or radiation, and for metastatic adenocarcinoma of the appendix or small bowel, and
  - No prior chemotherapy in the advanced setting, or
  - Previously not been suitable for therapy with bevacizumab but whose bevacizumab contraindication has since resolved, or
  - Received prior immunotherapy with UGI/VPEM or UGI/VPEM6 if MMR deficient metastatic colorectal adenocarcinoma, or
  - Received single agent capecitabine or fluorouracil treatment first-line as the result of frailty, but who are now well enough to receive combination chemotherapy, or
  - Progressed on single agent capecitabine or fluorouracil therapy first-line and treatment escalation/combination chemotherapy is desired

Patients should have:

- ECOG performance status less than or equal to 2
- Adequate marrow reserve, renal and liver function

Note: Consideration of first line oxaliplatin-based therapy (GICOXB) should be given for those patients who have Gilbert's Syndrome or who may be compromised by potential irinotecan toxicities

## EXCLUSIONS:

Patients must not have:

- Major surgery within 28 days of administration of therapy
- Untreated CNS metastases
- Severe renal impairment (Creatinine Clearance less than 30 mL/min)
- Suspected dihydropyrimidine dehydrogenase (DPD) deficiency (see Precautions)

## CAUTIONS:

- Patients with: 1) previous pelvic radiotherapy; 2) recent MI; 3) uncontrolled angina, hypertension, cardiac arrhythmias, congestive heart failure, 4) renal disease including proteinuria, 5) bleeding disorders, 6) previous anthracycline exposure, 7) prior radiation to the chest wall or other serious medical illness
- Patients with baseline greater than 3 loose BM per day (in patients without colostomy or ileostomy)
- Patients with recent (less than 6 months) arterial thromboembolic events
- Patients with baseline hyperbilirubinemia (greater than 26 micromol/L) not explained by degree of liver metastases

## TESTS AND MONITORING:

- **Baseline:** CBC and differential, Platelets, Creatinine, LFTs (Bilirubin, ALT, Alkaline Phosphatase), Albumin, sodium, potassium, DPYD test (not required if previously tested, or tolerated fluorouracil or capecitabine), dipstick or laboratory urinalysis for protein, Blood Pressure measurement and appropriate imaging study. Optional: CEA, CA 19-9.
- **Prior to each cycle:** CBC and differential, Platelets, Creatinine, LFTs (Bilirubin, ALT, Alkaline Phosphatase), Albumin, sodium, potassium, Blood Pressure measurement
  - **Prior to each even numbered cycles:** dipstick or laboratory urinalysis for protein
- 24 hour urine for protein if occurrence of proteinuria dipstick urinalysis shows 2+ or 3+ or laboratory urinalysis for protein is greater than or equal to 1g/L
- Blood Pressure measurement to be taken pre and post dose for first 3 cycles only and then pre-therapy with each subsequent visit.
- If clinically indicated: CEA, CA 19-9
- For patients on warfarin, weekly INR until stable warfarin dose established, then INR prior to each cycle.
- Quantitative evaluation of disease response status every six to twelve weeks; discontinue therapy if any progression of disease.
- Consider weekly nursing assessment for capecitabine toxicity in first two cycles and when increasing capecitabine dose.

## PREMEDICATIONS:

- Antiemetic protocol for high-moderate emetogenic chemotherapy with irinotecan, may not need any anti-emetic with capecitabine (see SCNAUSEA)
- Atropine may be required for treatment or prophylaxis of diarrhea (see precautions)
- Prochlorperazine should be avoided on the same day as irinotecan treatment due to the increased incidence of akathisia.

## TREATMENT:

A cycle equals -

Drug	Dose	BC Cancer Administration Guidelines
irinotecan	200 mg/m <sup>2</sup>	IV in 500 mL D5W over 1 hour 30 min
bevacizumab*	7.5 mg/kg*	IV in 100 mL NS over 15 minutes**
capecitabine <sup>†</sup>	800 mg/m <sup>2</sup> BID	PO x 14 days

Repeat every 21 days until disease progression.

**\*The bevacizumab dose should be recalculated for patients who experience more than a 10% change in body weight.**

**\*\* Observe for fever, chills, rash, pruritus, urticaria or angioedema and stop infusion and contact the physician if any of these occur. Infusion reactions should be treated according to severity. If the bevacizumab infusion is restarted then it should be given at an initial rate of 60 minutes or longer.**

**If acute hypertension (increase in BP measurement of greater than 20 mm Hg diastolic or greater than 160/100 if previously within normal limits) occurs during bevacizumab infusion – stop treatment. Resume at ½ the original rate of infusion if blood pressure returns to pretreatment range within one hour. If blood pressure does not return to pretreatment range within one hour – hold bevacizumab and subsequent infusions of bevacizumab should be given over 3 hours. Acute hypertension that is symptomatic (e.g. onset of headaches or change in level of consciousness) or BP measurement of greater than 180/110 that does not improve within one hour of stopping bevacizumab is an urgent situation that requires treatment.**

**Line should be flushed with Normal Saline pre and post dose as bevacizumab should not be mixed with dextrose solutions.**

**Capecitabine is available as 150 mg and 500 mg tablets (refer to Capecitabine Suggested Tablet Combination Table for dose rounding).**

- All patients should be advised to obtain an adequate supply of loperamide (IMODIUM®) with directions for the management of diarrhea.

## DOSE MODIFICATIONS

### Dose Levels for Toxicities

	Dose Level +1	Dose Level 0 (Starting Dose)	Dose Level -1	Dose Level -2
<b>irinotecan</b>	250 mg/m <sup>2</sup>	200 mg/m <sup>2</sup>	150 mg/m <sup>2</sup>	Discontinue Therapy
<b>capecitabine</b>	1000 mg/m <sup>2</sup>	800 mg/m <sup>2</sup>	500 mg/m <sup>2</sup>	Discontinue Therapy

**Patients who go through Cycle 1 at the Starting Dose without significant toxicity can be treated at Dose Level +1 on subsequent cycles.**

**DOSE MODIFICATIONS:**

**Capecitabine Dosing Based on DPYD Activity Score (DPYD-AS)**

Refer to "[Fluorouracil and Capecitabine Dosing Based on DPYD Activity Score \(DPYD-AS\)](http://www.bccancer.bc.ca/health-professionals/clinical-resources/cancer-drug-manual)" on [www.bccancer.bc.ca/health-professionals/clinical-resources/cancer-drug-manual](http://www.bccancer.bc.ca/health-professionals/clinical-resources/cancer-drug-manual).

**A. Dose Modifications for HEMATOLOGIC Toxicity**

Prior to a Cycle (Day 1)	Toxicity		Dose Level For Subsequent Cycles	
	Grade	ANC (x10 <sup>9</sup> /L)	irinotecan	capecitabine
<ul style="list-style-type: none"> <li>▪ If ANC less than 1.5 on Day 1 of cycle, hold treatment. Perform weekly CBC, maximum of 2 times.</li> <li>▪ If ANC is greater than or equal to 1.5 within 2 weeks, proceed with treatment at the dose level noted across from the <b>lowest ANC</b> result of the delayed week(s).</li> <li>▪ If ANC remains less than 1.5 after 2 weeks, discontinue treatment.</li> </ul>	1	greater than or equal to 1.5	Maintain dose level	Maintain dose level
	2	1.0 to less than 1.5	Maintain dose level	Maintain dose level
	3	0.5 to less than 1.0	↓ 1 dose level	↓ 1 dose level
	4	less than 0.5	↓ 2 dose levels	↓ 2 dose levels
	<b>Grade 4 neutropenia and greater than or equal to Grade 2 fever</b>		↓ 2 dose levels	↓ 2 dose levels

Prior to a Cycle (Day 1)	Toxicity		Dose Level For Subsequent Cycles	
	Grade	Platelets (x10 <sup>9</sup> /L)	irinotecan	capecitabine
<ul style="list-style-type: none"> <li>▪ If platelets less than 75 on Day 1 of cycle, hold treatment. Perform weekly CBC, maximum of 2 times.</li> <li>▪ If platelets greater than or equal to 75 within 2 weeks, proceed with treatment at the dose level noted across from the <b>lowest platelets</b> result of the delayed week(s).</li> <li>▪ If platelets remain less than 75 after 2 weeks, discontinue treatment.</li> </ul>	1	greater than or equal to 75	Maintain dose level	Maintain dose level
	2	50 to less than 75	Maintain dose level	Maintain dose level
	3	10 to less than 50	↓ 1 dose level	↓ 1 dose level
	4	less than 10	↓ 2 dose levels	↓ 2 dose levels

## B. Dose Modifications for NON-HEMATOLOGIC Toxicity

If Grade 2, 3 or 4 toxicities occur, daily administration of capecitabine should be immediately interrupted until these symptoms resolve or decrease in intensity to Grade 1.

Prior to a Cycle (Day 1)	Toxicity		Dose Level For Subsequent Cycles	
	Grade	Diarrhea	irinotecan	capecitabine
<ul style="list-style-type: none"> <li>▪ If diarrhea greater than or equal to Grade 2 on Day 1 of any cycle, hold treatment. Perform weekly checks, maximum 2 times.</li> <li>▪ If diarrhea less than Grade 2 within 2 weeks, proceed with treatment at the dose level noted across from the <b>highest</b> Grade experienced.</li> <li>▪ If diarrhea remains greater than or equal to Grade 2 after 2 weeks, discontinue treatment.</li> </ul>	1	Increase of 2 to 3 stools/day, or mild increase in loose watery colostomy output	Maintain dose level	Maintain dose level
	2	Increase of 4 to 6 stools, or nocturnal stools or mild increase in loose watery colostomy output	Maintain dose level	Maintain dose level
	3	Increase of 7 to 9 stools/day or incontinence, malabsorption; or severe increase in loose watery colostomy output	↓ 1 dose level	↓ 1 dose level
	4	Increase of 10 or more stools/day or grossly bloody colostomy output or loose watery colostomy output requiring parenteral support; dehydration	↓ 2 dose levels	↓ 2 dose levels

Prior to a Cycle (Day 1)	Toxicity		Dose Level For Subsequent Cycles	
	Grade	Stomatitis	irinotecan	capecitabine
<ul style="list-style-type: none"> <li>▪ If stomatitis greater than or equal to Grade 2 on Day 1 of any cycle, hold treatment. Perform weekly checks, maximum 2 times.</li> <li>▪ If stomatitis less than Grade 2 within 2 weeks, proceed with treatment at the dose level noted across from the <b>highest</b> Grade experienced.</li> <li>▪ If stomatitis remains greater than or equal to Grade 2 after 2 weeks, discontinue treatment.</li> </ul>	1	Painless ulcers, erythema or mild soreness	Maintain dose level	Maintain dose level
	2	Painful erythema, edema, or ulcers but can eat	Maintain dose level	Maintain dose level
	3	Painful erythema, edema, ulcers, and cannot eat	Maintain dose level	↓ 1 dose level
	4	As above but mucosal necrosis and/or requires enteral support, dehydration	Maintain dose level	↓ 2 dose levels

Prior to a Cycle (Day 1)	Toxicity		Dose Level For Subsequent Cycles	
	Grade	Palmar-Plantar Erythrodysesthesia (Hand-Foot Skin Reaction)	irinotecan	capecitabine
<ul style="list-style-type: none"> <li>▪ If hand-foot skin reaction greater than or equal to Grade 2 on Day 1 of any cycle, hold treatment. Perform weekly checks, maximum 2 times.</li> <li>▪ If hand-foot skin reaction less than Grade 2 within 2 weeks, proceed with treatment at the dose level noted across from the <b>highest</b> Grade experienced.</li> <li>▪ If hand-foot skin reaction remains greater than or equal to Grade 2 after 2 weeks, discontinue treatment.</li> </ul>	1	Skin changes (eg, numbness, dysesthesia, paresthesia, tingling, erythema) with discomfort not disrupting normal activities	Maintain dose level	Maintain dose level
	2	Skin changes (eg, erythema, swelling) with pain affecting activities of daily living	Maintain dose level	Maintain dose level
	3	Severe skin changes (eg, moist desquamation, ulceration, blistering) with pain, causing severe discomfort and inability to work or perform activities of daily living	Maintain dose level	↓ 1 dose level

**Renal dysfunction:**

Creatinine Clearance mL/min	Capecitabine Dose only
greater than or equal to 50	100%
30 to less than 50	75%
less than 30	Discontinue Therapy

**Cockcroft-Gault Equation:**

$$\text{Estimated creatinine clearance (mL/min)} = \frac{N \times (140 - \text{age in years}) \times \text{wt (kg)}}{\text{serum creatinine (micromol/L)}}$$

N = 1.23 male  
N = 1.04 female

**Proteinuria:**

There are 3 different measures of proteinuria that may be used to assess the need for modification of Bevacizumab therapy – urine dipstick analysis (measured in + values), laboratory urinalysis for protein (measured in g/L) and 24 hour urine collections for protein (measured in g/24 hours)

Urine dipstick analysis or laboratory urinalysis for protein should be performed at baseline and then prior to each even numbered cycle of therapy:

Degree of Proteinuria	
Neg or 1+ dipstick or less than 1 g/L laboratory urinalysis for protein	Administer bevacizumab dose as scheduled
2+ or 3+ dipstick or greater than or equal to 1 g/L laboratory urinalysis for protein	Administer bevacizumab dose as scheduled. Collect 24-hour urine for determination of total protein within 3 days before the next scheduled bevacizumab administration. <b>Adjust bevacizumab treatment based on the table below</b>
If urine dipstick shows 4+ at baseline or during treatment	Withhold bevacizumab and proceed with 24 hour urine collection.

24-Hour Urine Total Protein (G/24 hours)	Bevacizumab Dose
less than or equal to 2	100%
greater than 2 to 4	Hold dose and recheck 24 hour urine every 2 weeks, resume therapy when less than or equal to 2g/24 hour
greater than 4	Discontinue Therapy

**Hypertension:**

Blood Pressure (mm Hg)	Bevacizumab Dose
less than or equal to 160/100	100%
greater than 160/100 asymptomatic	100% Notify physician and start or adjust antihypertensive therapy*
Hypertensive Crisis	Discontinue Therapy

- **Antihypertensive therapy may include hydroCHLORothiazide 12.5 to 25 mg PO once daily, ramipril (ALTACE®) 2.5 to 5 mg PO once daily, or amlodipine (NORVASC™) 5 to 10 mg PO once daily.**

## PRECAUTIONS:

- Diarrhea:** may be life threatening and requires prompt, aggressive treatment.
  - Early diarrhea** or abdominal cramps occurring within the first 24 hours is treated with **atropine** 0.3 to 1.2 mg IV or SC. Prophylactic atropine may be required for subsequent treatments.
  - Late diarrhea** has an onset of 5 to 11 days post-treatment, a duration of 3 to 7 days and must be treated promptly with **loperamide** (eg, IMODIUM®). The loperamide dose is higher than recommended by the manufacturer. Instruct patient to have loperamide on hand and start treatment at the first poorly formed or loose stool, or earliest onset of more frequent stool than usual:
    - 4 mg stat**
    - then 2 mg every 2 hours until diarrhea-free for 12 hours**
    - may take 4 mg every 4 hours at night
  - The use of drinks such as Gatorade® or Powerade® to replace fluid & body salts is recommended.
  - Consideration should be given to the use of an oral fluoroquinolone (e.g., ciprofloxacin) in patients with persistent diarrhea despite adequate loperamide or if a fever develops in the setting of diarrhea, even without neutropenia. If diarrhea persists for longer than 48 hours then hospitalization for parenteral hydration should be considered.
- Other cholinergic symptoms:** may occur during or shortly after infusion of irinotecan including rhinorrhea, increased salivation, lacrimation, diaphoresis and flushing. These should be treated with atropine 0.3 mg – 0.6 mg IV or SC. This dose may be repeated at the physician's discretion. Blood pressure and heart rate should be monitored. Prophylactic atropine may be required for subsequent treatments.
- Neutropenia:** Fever or other evidence of infection must be assessed promptly and treated aggressively.
- Gastrointestinal perforations and wound dehiscence:** Can be fatal. Typical presentation is reported as abdominal pain associated with symptoms such as constipation and vomiting. Bevacizumab should be discontinued in patients with gastrointestinal perforation or wound dehiscence requiring medical intervention.
- Hemorrhage:** Bevacizumab has been associated with hemorrhage. Cases of CNS hemorrhage, some with fatal outcome, have been observed. Patients should be monitored for signs and symptoms of CNS bleeding. If Grade 3/4 hemorrhage occurs, discontinue Bevacizumab. Patients with significant bleeding diatheses should not receive Bevacizumab. Platelet inhibitory medications such as NSAIDs (including ASA at doses greater than 325 mg/day) should be discontinued prior to institution of Bevacizumab. COX-2 inhibitors are permissible. **For patients on warfarin, see Thrombosis (for bevacizumab) and Drug Interactions (for capecitabine).**
- Thrombosis:** A history of arterial thromboembolic events or age greater than 65 years is associated with an increased risk of arterial thromboembolic events with Bevacizumab. If Grade 3 thromboembolic event or incidentally discovered pulmonary embolus arises, hold Bevacizumab for 2 weeks, then consider resumption of Bevacizumab if risks of tumour-related hemorrhage are judged low AND the patient is on a stable dose of anticoagulant. If a second Grade 3 thrombosis occurs, or if a Grade 4 thrombosis occurs, discontinue Bevacizumab. Patients on warfarin should have INR checked frequently, at least once per cycle, while receiving Bevacizumab. In patients on warfarin with an elevated INR, it is recommended to **hold the bevacizumab if INR is greater than 3.0.**
- Proteinuria:** Has been seen in all clinical trials with Bevacizumab to date and is likely dose-dependent. If proteinuria of greater than or equal to 2g/24 hr persists for more than 3 months, consider further investigations - possibly a renal biopsy.
- Hypertension:** Has been seen in all clinical trials with Bevacizumab to date and is likely dose-dependent. The most commonly used therapies are Calcium Channel Blockers, ACE Inhibitors and Diuretics. Blood pressure should be monitored through routine vital signs evaluations. If hypertension is poorly controlled with adequate medication, discontinue Bevacizumab.
- Reversible Posterior Leukoencephalopathy Syndrome:** Rarely, patients receiving bevacizumab may develop seizures, headache, altered mental status, visual disturbances, with or without associated hypertension consistent with RPLS. May be reversible if recognized and treated promptly.



9. **Congestive Heart Failure:** Has been reported in up to 3.5% of patients treated with bevacizumab. Most patients showed improvement in symptoms and/or LVEF following appropriate medical therapy.
10. **Myocardial ischemia and angina occurs rarely in patients receiving fluorouracil or capecitabine.** Development of cardiac symptoms including signs suggestive of ischemia or of cardiac arrhythmia is an indication to discontinue treatment. If there is development of cardiac symptoms patients should have urgent cardiac assessment. Generally re-challenge with either fluorouracil or capecitabine is not recommended as symptoms potentially have a high likelihood of recurrence which can be severe or even fatal. Seeking opinion from cardiologists and oncologists with expert knowledge about fluorouracil / capecitabine toxicity is strongly advised under these circumstances. The toxicity should also be noted in the patient's allergy profile.
11. **Gilbert's syndrome:** Increases the risk of irinotecan-induced toxicity. A screen for Gilbert's Syndrome using direct/indirect serum bilirubin is recommended.
12. **Hepatic dysfunction:** Irinotecan has not been studied in patients with bilirubin greater than 35 micromol/L or ALT greater than 3x the upper limit of normal if no liver metastases, or ALT greater than 5x the upper limit of normal with liver metastases. The risk of severe neutropenia may be increased in patients with a serum bilirubin of 17 to 35 micromol/L.
13. **Pulmonary toxicity:** Severe pulmonary toxicity consisting of dyspnea, fever and reticulonodular pattern on chest x-ray has been reported rarely. Supportive care is required.
14. **Prior pelvic radiotherapy** or radiotherapy to greater than 15% of the bone marrow bearing area may increase the degree of myelosuppression associated with this regimen, and caution is recommended in these cases. Close monitoring of the CBC is essential.
15. **Potential Drug Interactions:** Anticonvulsants and other drugs which induce Cytochrome P450 3A4 isoenzyme activity e.g. Carbamazepine, Phenytoin and St John's Wort may decrease the therapeutic and toxic effects of **irinotecan**. Prochlorperazine may increase the incidence of akathisia and should be avoided on the day of **irinotecan** treatment.
16. **Possible drug interaction with capecitabine and warfarin** has been reported and may occur at any time. For patients on warfarin, weekly INR during capecitabine therapy is recommended until a stable warfarin dose is established. Thereafter, INR prior to each cycle. Consultation to cardiology/internal medicine should be considered if difficulty in establishing a stable warfarin dose is encountered. Upon discontinuation of capecitabine, repeat INR weekly for one month.
17. **Possible drug interaction with capecitabine and phenytoin and fosphenytoin** has been reported and may occur at any time. Close monitoring is recommended. Capecitabine may increase the serum concentration of these two agents.
18. **Dihydropyrimidine dehydrogenase (DPD) deficiency** may result in severe and unexpected toxicity – stomatitis, diarrhea, neutropenia, neurotoxicity – secondary to reduced drug metabolism. This deficiency is thought to be present in about 3% of the population.

**Call the GI Systemic Therapy physician at your regional cancer centre or the GI Systemic Therapy Chair Dr. Theresa Chan at (604) 930-2098 with any problems or questions regarding this treatment program.**

#### REFERENCES:

1. Patt YZ, et al. Capecitabine plus irinotecan: A highly active first-line treatment for metastatic colorectal cancer. Proc Am Soc Clin Oncol 2003;22:281a (abstract 1130).
2. Kerr DJ, et al. A phase I/II study of CPT-11 in combination with capecitabine as first line chemotherapy for metastatic colorectal cancer. Proc Am Soc Clin Oncol 2002;21:100a (abstract 643).
3. Delord P, et al. Dose escalation and pharmacokinetic study of capecitabine (Xeloda) and irinotecan (CPT-11) in gastrointestinal tumors: preliminary results. Proc Am Soc Clin Oncol 2002;21:100a (abstract 397).
4. Bajetta E, et al. Randomized multicenter Phase II trial of two different schedules of irinotecan combined with capecitabine as first-line treatment in metastatic colorectal carcinoma. Cancer 2004;100(2):279-87.
5. Wasserman E, et al. Severe CPT-11 toxicity in patients with Gilbert's syndrome: two case reports. Ann Oncol 1997;8(10):1049-51.
6. Mathijssen RHJ, et al. Effects of St. John's Wort on irinotecan metabolism. J Natl Cancer Inst 2002;94(16):1247-9.
7. Hurwitz H et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. NEJM 2004; 350(23):2335-42.

8. Kabbinavar F, et al. Addition of Bevacizumab to bolus Fluorouracil and Leucovorin in first-line metastatic colorectal cancer: results of a randomized phase II trial. *J Clin Oncol* 2005;23:3697-705.
9. Kabbinavar F, et al. Combined analysis of efficacy: the addition of bevacizumab to fluorouracil/leucovorin improves survival for patients with metastatic colorectal cancer. *J Clin Oncol* 2005;23:3706-12.
10. Saltz LB, et al. Simplification of bevacizumab (bev) administration: Do we need 90, 60, or even 30 minute infusion times? *J Clin Oncol (Meeting Abstracts)* 2006;24(18\_suppl):3542.