



BC Cancer Colon Screening 2017 Program Results

Date Published: September 2019

Table of Contents

Program Overview	3
<hr/>	
Program Results	5
Program Uptake	5
FIT Results	11
Colonoscopy Results	12
Wait Times	17
Quality Assurance	18
Summary	18
<hr/>	
Appendix: Performance Indicator Glossary	19

PROGRAM OVERVIEW

Colon cancer screening in B.C. is organized under a partnership framework with regional health authorities, laboratory service providers, primary care providers and specialists. BC Cancer provides oversight for organized cancer screening in B.C., and supports:

- development of provincial policies, guidelines and standards,
- strategies to increase public and health care provider awareness, including both benefits and limitations of screening,
- correspondence to eligible British Columbians about results, follow-up and rescreening,
- quality assurance and quality improvement, and
- reporting and monitoring of system performance and screening outcomes.

In B.C., regional health authorities are responsible for the planning and delivery of healthcare services within their geographic areas. Health Authorities and community health service providers work with BC Cancer Screening to provide high quality screening and diagnostic services.

Primary care providers play the important role of identifying eligible individuals for screening. BC Cancer provides material to help primary care providers discuss the benefits and limitations of screening with their patients. Once the decision to screen is made, the primary care provider directs the patient to the appropriate screening test, and supports them throughout their screening journey.

In addition, as part of the Indigenous Cancer Strategy, BC Cancer Screening works collaboratively with the First Nations Health Authority, Métis Nation British Columbia and the B.C. Association of Aboriginal Friendship Centres to improve cancer screening access and participation of Indigenous people.


At this time Northern Health Authority follows their own colon screening processes for referral and recall and does not provide data to the Provincial program. Therefore, no monitoring of the efficacy and quality of colon screening can be done for the people living in the area comprising the Northern Health Authority.

The Colon Screening Program started in B.C. in November 2013. The data provided in this report is based on screening results for British Columbians registered in the Colon Screening Program.

The Screening Process

The screening pathway is initiated by primary care providers referring asymptomatic individuals 50 to 74 years of age for a screening test – either the fecal immunochemical test (FIT) or colonoscopy, depending on the patient’s risk of developing colorectal cancer. Figure 1 provides an overview of the colon screening process.

FIGURE 1: COLON SCREENING PROCESS OVERVIEW



PROGRAM RESULTS

In order to prevent inappropriate disclosure of health-related information, all integers presented in this report have been randomly rounded up or down to the nearest five using Statistics Canada methodology.

A fecal immunochemical test (FIT) is the screening test offered for average risk individuals in BC. Through the spring of 2017 the proportion of abnormal tests began increasing. The laboratory providers in BC worked with the manufacturer and identified some differences in reagent production. The abnormal test rate continued to increase in August and September 2017 and a decision was made by the laboratories to suspend FIT in BC in early October 2017 while working with the manufacturer regarding the reagent. FIT was reinstated mid-December 2017 after the laboratories completed quality evaluations. The effect of the reagent changes and the unavailability of the test in BC for 2 ½ months in 2017 can be seen below with regard to volume, participation, abnormal FIT rate and positive predictive values.

a) Program Uptake


Asymptomatic British Columbians, ages 50 to 74, can enter into the Colon Screening Program by visiting their primary care provider. The primary care provider assesses the individual's risk of developing colorectal cancer and orders the appropriate screening test –FIT for an average risk individual and colonoscopy for higher than average risk.

Primary care providers enroll asymptomatic average risk individuals by selecting the appropriate option on the laboratory requisition form. Colonoscopy referral for higher than average risk individuals is sent directly to the Colon Screening Program.

Figure 2 shows the volume of FIT results received by the Colon Screening Program by quarter since the inception of the provincial program. There continues to be a higher proportion of first time screeners registering in the program. The number of people returning for subsequent rounds of screening is growing as expected. Volumes for the 4th quarter in 2017 are low due to the FIT suspension that occurred during this quarter. The proportion of FITs with results copied to the Colon Screening Program increased for the first three quarters of 2017 up to 77.5% (Figure 3).

In 2017, 24% of patients had a repeat FIT within 21 months following a negative FIT in the program. Early return to screening does not increase the uptake of colon screening in B.C. but utilizes screening resources.


FIGURE 2: NUMBER OF FIT RESULTS RECEIVED BY THE COLON SCREENING PROGRAM OVER TIME



NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. Integers have been rounded as per Statistics Canada methodology.
3. FIT was unavailable in B.C. for most of Q4 2017.

FIGURE 3: PROPORTION OF FITs REGISTERED WITH THE COLON SCREENING PROGRAM FOR BRITISH COLUMBIANS AGES 50-74




NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. An Individual may have multiple FITs performed in any time period.
3. FIT was unavailable in B.C. for most of Q4 2017.

Figure 4 and Figure 5 demonstrate that the number of colonoscopies performed in individuals at higher than average risk has continued to increase. This includes participants with a high risk family history defined as one first degree relative (i.e. parent, sibling or child) with colorectal cancer diagnosed under the age of 60 or two or more first degree relatives with colorectal cancer diagnosed at any age. A high risk family history is the colonoscopy indication in 30% of higher than average risk referrals while a personal history of adenoma(s) accounts for 70% of higher than average risk patients referred to Health Authorities for colonoscopy in 2017.


FIGURE 4: NUMBER OF REFERRALS FOR PATIENTS WITH PERSONAL HISTORY OF ADENOMA



NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. An individual may have multiple referrals.
3. Integers have been rounded as per Statistics Canada methodology.

FIGURE 5: NUMBER OF REFERRALS FOR PATIENTS WITH FAMILY HISTORY OF COLON CANCER




NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. An individual may have multiple referrals.
3. Integers have been rounded as per Statistics Canada methodology.

In 2017, the program received 192,585 FIT results on 189,873 British Columbians ages 50 to 74, and 8,789 colonoscopies were completed for higher than average risk individuals. 30.6% of the age eligible population has had a FIT within the Colon Screening Program in the past 30 months (Figure 5). Of these, 52% were female and the mean age of individuals was 62 years.

Figure 6 shows FIT participation by age and sex. Regional variation is shown in Figure 7. This does not account for higher than average risk participants undergoing colonoscopy or those participants up to date with screening (ie. Previous abnormal FIT with a normal colonoscopy to be recalled for FIT 10 years following colonoscopy).


FIGURE 6: PROGRAM PARTICIPATION RATE IN B.C. BY AGE AND SEX



NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. Population data source: P.E.O.P.L.E 2018 (Sept 2018), BC STATS, Service BC, BC Ministry of Citizen’s Services

FIGURE 7: PROGRAM PARTICIPATION RATE BY HEALTH AUTHORITY



NOTES:


1. Colon Screening Program data extraction date: 04/02/2019.
2. Population data source: P.E.O.P.L.E 2018 (Sept 2018), BC STATS, Service BC, BC Ministry of Citizen’s Services

The following sections describe the Colon Screening Program results from January 1, 2017 to December 31, 2017.

b) FIT Results

The percent of FIT results that were abnormal in 2017 was 17.6%, this is an increase from 13.5% in 2016 and most likely related to the FIT manufacturer reagent as opposed to population differences. Figure 8 demonstrates that abnormal FIT results were more common in males and increase with age, which reflects the prevalence of colorectal cancer.

FIGURE 8: FIT POSITIVITY BY AGE GROUP AND SEX



NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.


c) Colonoscopy Results

Participants with Abnormal FIT Results

In 2017, a total of 34,540 program participants with abnormal FIT results were referred to regional health authorities for colonoscopy assessment. After initial assessment by health authority staff, 71.1% proceeded to have a colonoscopy with outcome data captured by the Colon Screening Program, 10.4% declined colonoscopy or were unable to be contacted, 9.0% were deemed ineligible for the program and 9.5% did not proceed to colonoscopy through the program but likely obtained follow-up through a provider directly. This underscores the importance of having primary care providers assess a potential participant’s understanding that an abnormal FIT result requires a colonoscopy to complete the screening episode. This assessment should occur prior to proceeding with FIT. Further education is required regarding screening eligibility.

Figure 9 summarizes the outcomes for those with abnormal FIT results. Of the 22,410 cases with available pathology information 53.3% were found to have colorectal cancer or a pre-cancerous polyp: 340 (1.6%) cases for whom a colorectal cancer was found, 3,400 (15.2%) cases with high risk polyp(s) identified, 1,525 (6.8%) cases with multiple (3 or more) low risk polyps and 6,650 (29.7%) cases with 1 or 2 low risk polyp(s). For the cancers, 115 (34.8%) were located on the left side of the colon, 105 (31.8%) were right-sided and 105 (31.8%) were in the rectum.

FIGURE 9: COLONOSCOPY FINDINGS FOR THOSE WITH AN ABNORMAL FIT RESULT



Quality indicators help assess the effectiveness of the colonoscopy. These include cecal intubation and adequate bowel preparation. The unadjusted cecal intubation rate was 98.1% and the adequate bowel preparation rate was 98.2% in colonoscopies done for patients with abnormal FIT results.

The positive predictive value (PPV) of a test is a measure of performance. It represents the proportion of individuals with an abnormal FIT who have cancer or pre-cancerous polyps at follow-up colonoscopy. The PPV of FIT is presented in Table 1. For ages 50 to 74 combined, the PPV for any neoplasia (cancer and any pre-cancerous polyp) is 53.2% while the PPV for colorectal cancer and high risk polyps is 16.7%. The PPV of FIT increases with age and is higher in males than females.

TABLE 1: POSITIVE PREDICTIVE VALUE OF THE FIT

	Cancer	High Risk Polyp(s)	Multiple Low Risk Polyps	Low risk polyp	Any Neoplasia
All	340 (1.5%)	3,400 (15.2%)	1,525 (6.8%)	6,650 (29.7%)	11,915 (53.2%)
By FIT					
First FIT	240 (1.8%)	2,295 (16.8%)	925 (6.8%)	3,890 (28.5%)	7,345 (53.8%)
Subsequent FIT	100 (1.1%)	1,100 (12.6%)	605 (6.9%)	2,755 (31.5%)	4,560 (52.1%)
By Sex					
Females	140 (1.4%)	1,220 (11.8%)	475 (4.6%)	2,865 (27.7%)	4,700 (45.5%)
Males	195 (1.6%)	2,180 (18.1%)	1,050 (8.7%)	3,780 (31.3%)	7,205 (59.7%)
By Age group					
50-54	40 (1.0%)	500 (12.0%)	185 (4.5%)	1,090 (26.2%)	1,820 (43.8%)
55-59	55 (1.2%)	665 (14.1%)	240 (5.1%)	1,425 (30.2%)	2,380 (50.4%)
60-64	70 (1.4%)	805 (16.0%)	330 (6.6%)	1,490 (29.7%)	2,690 (53.5%)
65-69	85 (1.7%)	825 (16.7%)	420 (8.5%)	1,510 (30.6%)	2,835 (57.4%)
70-74	90 (2.5%)	605 (17.0%)	355 (9.9%)	1,130 (31.7%)	2,180 (61.2%)

NOTES:


1. Colon Screening Program data extraction date: 04/02/2019.
2. Integers have been rounded as per Statistics Canada methodology.

Higher than Average Risk Participants with Personal History of Adenomas

During the report period, 6,600 referrals for colonoscopy assessment were sent to the Health Authorities for higher than average risk screening due to a personal history of adenomas. After initial assessment by health authority staff, 70.3% proceeded to have a colonoscopy with outcome data captured by the Colon Screening Program, 7.9% declined colonoscopy or were unable to be contacted, 11.0% were deemed ineligible for the program and 10.9% did not proceed to colonoscopy through the program but likely obtained follow-up through a provider directly. This emphasizes the need for ongoing primary care education regarding program eligibility.

Figure 10 summarizes colonoscopy findings for those with a personal history of adenomas. Of the 4,535 cases with available follow-up information, 64.8% were found to have colorectal cancer or a precancerous polyp.

FIGURE 10: COLONOSCOPY FINDINGS FOR THOSE WITH A PERSONAL HISTORY OF ADENOMAS



NOTES:


1. Colon Screening Program data extraction date: 04/02/2019
2. High risk polyp(s) include the following: tubular adenoma or sessile serrated adenoma/polyp ≥ 10 mm in diameter, villous or tubulovillous adenoma, sessile serrated adenoma/polyp with dysplasia and traditional serrated adenoma.
3. Integers have been rounded as per Statistics Canada methodology.

Higher than Average Risk Participants with Family History of Colon Cancer

During the report period, 1,720 referrals for pre-colonoscopy assessment were sent to the Health Authorities for those with a family history of colon cancer. After initial assessment by health authority staff, 69.6% proceeded to have a colonoscopy with outcome data captured by the Colon Screening Program, 6.1% declined colonoscopy or were unable to be contacted, 19.0% were deemed ineligible for the program and 5.3% did not proceed to colonoscopy through the program but likely obtained follow-up through a provider directly. This emphasizes the need for ongoing primary care education on the eligibility for screening.

Figure 11 summarizes colonoscopy findings for higher risk participants with a family history of colon cancer. Of the 1,175 cases with available follow-up information, 48.3% were found to have colorectal cancer or a precancerous polyp.

FIGURE 11: COLONOSCOPY FINDINGS FOR THOSE WITH A FAMILY HISTORY



NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. High risk polyp(s) include the following: tubular adenoma or sessile serrated adenoma/polyp ≥ 10 mm in diameter, villous or tubulovillous adenoma, sessile serrated adenoma/polyp with dysplasia and traditional serrated adenoma.
3. Integers have been rounded as per Statistics Canada methodology.

In the higher than average risk patients undergoing colonoscopy, the unadjusted cecal intubation rate was 98.0%, and 97.7% had an adequate bowel preparation.

Detection of neoplasia in screening colonoscopy for those with a personal history and those with a family history of adenomas are presented in Tables 2 and 3. Detection of cancer and high risk polyps in the higher than average risk groups is lower than observed for those with an abnormal FIT result.

TABLE 2: DETECTION OF NEOPLASIA IN SCREENING COLONOSCOPY FOR THOSE WITH A PERSONAL HISTORY OF ADENOMAS

Age Group	Cancer	High Risk Polyp(s)	Multiple		Any Neoplasia
			Low Risk Polyps	Low risk polyp	
All	20 (0.4%)	615 (13.6%)	590 (13.0%)	1715 (37.8%)	2,940 (64.8%)
By Sex					
Females	10 (0.6%)	205 (12.2%)	170 (10.1%)	610 (36.3%)	980 (58.3%)
Males	10 (0.4%)	420 (14.7%)	425 (14.9%)	1110 (38.9%)	1,960 (68.8%)
By Age group					
50-54	0 (0.0%)	30 (10.3%)	30 (10.5%)	100 (35.1%)	160 (56.1%)
55-59	0 (0.0%)	95 (13.2%)	75 (10.4%)	245 (34.0%)	415 (57.6%)
60-64	0 (0.0%)	135 (12.6%)	130 (12.1%)	415 (38.8%)	690 (64.2%)
65-69	5 (0.4%)	195 (14.4%)	190 (14.0%)	540 (39.9%)	925 (68.3%)
70-74	5 (0.5%)	165 (15.0%)	165 (14.9%)	415 (37.7%)	755 (68.6%)

NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. Integers have been rounded as per Statistics Canada methodology.

TABLE 3: DETECTION OF NEOPLASIA IN SCREENING COLONOSCOPY FOR THOSE WITH A FAMILY HISTORY OF COLORECTAL CANCER

Age Group	Cancer	High Risk Polyp(s)	Multiple		Any Neoplasia
			Low Risk Polyps	Low risk polyp	
All	5 (0.4%)	110 (9.4%)	80 (6.8%)	370 (31.5%)	565 (48.3%)
By Sex					
Females	0 (0.0%)	70 (10.7%)	30 (4.6%)	175 (26.7%)	270 (41.2%)
Males	5 (1.0%)	45 (8.7%)	50 (9.7%)	195 (37.9%)	295 (57.3%)
By Age group					
50-54	0 (0.0%)	10 (5.0%)	5 (2.4%)	60 (30.0%)	75 (37.5%)
55-59	0 (0.0%)	30 (9.7%)	20 (6.5%)	90 (29.0%)	140 (45.2%)
60-64	5 (1.9%)	20 (7.4%)	20 (7.4%)	90 (32.7%)	130 (48.1%)
65-69	0 (0.0%)	25 (11.4%)	20 (9.1%)	80 (36.4%)	125 (56.8%)
70-74	0 (0.0%)	20 (11.8%)	15 (9.1%)	55 (32.4%)	95 (57.6%)

NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. Integers have been rounded as per Statistics Canada methodology.

Table 4 compares detection rates for the three different populations participating in B.C.’s Colon Screening Program.

TABLE 4: DETECTION RATE BY POPULATION TYPE

Pathology	Personal History of		
	FIT Positive	Adenoma	Family History
Total	22,410	4,535	1,175
Cancer	340 (1.5%)	20 (0.4%)	5 (0.4%)
High Risk Polyp	3,400 (15.2%)	615 (13.6%)	110 (9.4%)
Any Neoplasia	11,915 (53.2%)	2940 (64.8%)	565 (48.3%)
No Neoplasia	10,505 (46.9%)	1,595 (35.2%)	600 (51.5%)


NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.
2. Any neoplasia includes high risk polyps, multiple low risk polyps and low risk polyps.
3. No neoplasia includes patients where no specimens were taken at colonoscopy and other polyps/specimens being removed.
4. Integers have been rounded as per Statistics Canada methodology.

d) Wait Times

Wait times for colonoscopy after an abnormal FIT result are shown in 6-month intervals in Figure 12. The target time from an abnormal FIT result to colonoscopy is 60 days. The rapid increase in the number of individuals requiring colonoscopy, since the launch of the program in November 2013, has created a challenge in every health authority to meet the 60 day target. The increase in the proportion of abnormal FIT results also increased demand for colonoscopy in 2017. It is recognized that there are many other indications for endoscopy services.

FIGURE 12: WAIT TIME FROM ABNORMAL FIT TO COLONOSCOPY



NOTES:

1. Colon Screening Program data extraction date: 04/02/2019.

e) Quality Assurance

All colonoscopists providing procedures for Colon Screening Program participants in B.C. are encouraged to participate in direct observation of procedural skills (DOPS). DOPS is a formative assessment of a physician's performance of colonoscopy in terms of technical skill as well as patient and staff interaction. The DOPS process involves two trained assessors simultaneously and independently observing a physician perform two consecutive colonoscopies and completing a validated form. The assessors provide constructive feedback to the physician in written and verbal formats.

All endoscopy units providing procedures for Colon Screening Program participants in B.C. are expected to participate in the Global Rating Scale-Canada (GRS-C). GRS-C is a biannual survey to assess all aspects of endoscopic quality assurance at the level of the endoscopy unit. The survey is a patient-centered tool which enables units to identify areas not yet meeting quality standards and design action plans for quality improvement. The survey exists on a web-based platform supported by the Canadian Association of Gastroenterology.

Annual quality reports are sent to health authorities, primary care providers, colonoscopists and pathologists participating in the program with individual and aggregate performance statistics.

e) Summary

The following are some key findings based on the 2017 data:

- Due to FIT being suspended for part of 2017, participation has decreased by 3.0% between 2016 and 2017. The participation rate calculation in this report does not take in to account age eligible British Columbians who may be up to date with colon screening but who's results are not captured by the program.
- The number needed to screen to detect one cancer is 535.
- The number needed to screen to detect one cancer or high risk polyp is 44.
- The number of participants with an abnormal FIT needed to undergo colonoscopy to detect one cancer is 60.
- The number of participants with an abnormal FIT needed to undergo colonoscopy to detect one cancer or high risk polyp is 5.
- 60% of colonoscopists completed DOPS.
- There are further opportunities to support primary care providers in using the Colon Screening Program:
 - 24% of patients are having FIT ordered less than 21 months from the last negative FIT.
 - 10% of patients being referred for colonoscopy are assessed by the Health Authority staff to be ineligible for the program (colonoscopy in the last 5 years, personal history of CRC, incorrect family history or medically unfit).
 - 10% of patients being referred for colonoscopy decline or do not respond when contacted.

APPENDIX – PERFORMANCE INDICATOR GLOSSARY

Program Participation Rate

Percentage of British Columbia screen-eligible population, ages 50-74, who completed a fecal immunochemical test (FIT) registered with the Colon Screening Program within a 30-month period. Prevalence adjusted participation is used, as individuals who have had a previous colorectal cancer diagnosis at any point in time are no longer eligible to participate in the Colon Screening Program, and are therefore excluded from the population estimate.

$$\text{Program Participation rate} = \frac{\text{Number of patients with a successful FIT referral}}{\text{Prevalence adjusted BC population as of December 2017}} \times 100$$

FIT Positivity Rate

FIT positivity rate is defined as the number of satisfactory FITs with an abnormal result.

$$\text{FIT Positivity Rate} = \frac{\text{Number of FITs with an abnormal result}}{\text{Number of satisfactory FITs}} \times 100$$

FIT Positive Predicted Value (PPV)

FIT positive predicted value is defined as the proportion of satisfactory FITs resulting in pathological confirmation, where pathology result is some specified category of neoplasia.

$$\text{FIT PPV} = \frac{\text{Number of satisfactory FITs with pathologically confirmed neoplasia}}{\text{Number of satisfactory FITs with diagnostic data confirmation}} \times 100$$

Detection of Neoplasia (Higher Than Average Risk Patients)

Neoplasia detection rates defined as the proportion of colonoscopy procedures resulting in pathological confirmation, where the pathology result is some specified category of neoplasia.

$$\text{Neoplasia Detection Rate} = \frac{\text{Number of colonoscopies with pathologically confirmed neoplasia}}{\text{Number of colonoscopies}}$$

Cecal Intubation Rate (Unadjusted)

Unadjusted cecal intubation rate is defined as proportion of colonoscopy procedures in which the cecum was intubated.

$$\text{Unadjusted Cecal Intubation Rate} = \frac{\text{Number of procedures w/ cecal intubation}}{\text{Total number of colonoscopies}} \times 100$$

Adequate Bowel Preparation Rate

Adequate bowel preparation rate is defined as the proportion of colonoscopy procedures where the bowel preparation was defined as either 'excellent', 'good', or 'fair' (i.e. not 'poor').

$$\text{Adequate Bowel Preparation Rate} = \frac{\text{Number of colonoscopy procedures w/ adequate bowel prep}}{\text{Total number of colonoscopies}} \times 100$$

Wait Time to Follow-Up Colonoscopy

Wait time to follow-up colonoscopy is defined as the number of days elapsed between an abnormal FIT result and date of follow-up colonoscopy, for patients who had an abnormal FIT result and have received a colonoscopy.